

# **Severe Allergy Alert Form**

# Part 1

This form must be completed when the student is first registered or re-registered with the Holy Spirit School Division, or when the student's allergies change.

STUDENT INFORMATION (To be completed by Parent or Guardian)		
Name of Student:	Date of Birth:	
Address:		
Home Telephone:	Medic Alert I.D.:	
Name of Parent/Guardian:	Daytime #:	
Emergency Contact Person:	Daytime #:	
Emergency Contact Person	Daytime #:	
PHYSICIAN INFORMATION (To be completed by Physician)		
Nature of Allergy/Allergens:		
Symptoms of Reaction:		
Recommended Response to Reaction:		
Medication:	Dosage:	
Additional Instructions or Information:		
Name of Physician:	Telephone #:	
Signature of Physician:	Date:	
The personal information on this form is collected under the authority of the School Act and The purpose of this collection is to respond to potential emergency situations involving your slife-threatening allergy. If you have any questions concerning the collection, use or disclosule either in writing or by telephone.	student whom you have identified as subject to a potentially	



# **Severe Allergy Alert Form**

Part 2

This form must be completed by a parent/legal guardian or independent student when a student's attendance at school is affected by a dangerous, life-threatening allergy. The information gathered in this form must be reviewed (and confirmed and updated) annually or sooner if the student's condition changes.

		1
Student's Name:		
ALLERGY - DESCRIPTION This student has a DANGEROUS, life-threatening allergy to the following	<b>j</b> :	
And all substances containing them in any form or amount, including the following kinds of items:		Place Student's Photo Here
AVOIDANCE The key to preventing an emergency is ABSOLUTE AVOIDANCE of these	se allergens at all times.	
GENERAL PRECAUTIONS		
If assistance with administering medication may be required, o	complete Request for Assistance to	o Administer Medication Form.
SYMPTOMS FOLLOWING EXPOSURE TO A PARTICUAL	LR MATERIAL CAN INCLUDE:	
<ul> <li>Hives and itchiness on any part of the body;</li> <li>Nausea, vomiting, diarrhea;</li> <li>Difficulty breathing or swallowing</li> <li>Panic or sense of doom;</li> <li>Throat tightness or closing</li> </ul>	<ul> <li>Swelling of any body parts, especially eyelids, lips, face or tongue;</li> <li>Coughing, wheezing or change of voice;</li> <li>Fainting or loss of consciousness;</li> <li>Other (please specify)</li> </ul>	
EMERGENCY MEASURES		
Get EpiPen (epinephrine) or other Medication and administ HAVE SOMEONE CALL AN AMBULANCE and advise that a Unless the student is resisting, lay student down, tilt head bace Cover and reassure student. Record the time at which EpiPen (epinephrine) was adminis Have someone call the parent. If the ambulance has not arrived in 10 – 15 minutes, and breate Even if symptoms subside, students require medical attention ambulance. Have the parent / guardian or a school staff member accompated Provide ambulance and / or hospital personnel with a copy of or Medication was administered.  I understand why I have been asked to disclose the above student's ider to the disclosure. I voluntarily give the school consent to place a copy of in appropriate locations within the school, take the Emergency Measures	a child is having an anaphylactic reaction is and elevate legs.  Itered.  Ithing difficulties are present, administer a because there may be a delayed reaction any the student to the hospital. Ithe Severe Allergy Alert Form for the studentifying information and I am aware of the this form in the student's cumulative student and share this information, as necessare.	a second EpiPen (epinephrine). on. Take the student to hospital immediately in the udent and the time at which the EpiPen (epinephrine) erisks or benefits of consenting or refusing to consent ident record, post this form including student's picture ry, with the staff of the school and health providers.
	ignature of Parent/Guardian or Independ	·



# Severe Allergy Resource Package Sample Letter to Parents

#### Dear Parent

I am writing to you on behalf of one of our students and parents. The student has a potentially life-threatening allergy to << insert the name of the allergen >>.

<<p><<If peanut butter or even the tiniest amount of peanut, a peanut product such as peanut oil, or any type of nut enters the student's body through the eyes, nose, or mouth, the student experiences very strong reactions. The student's face swells and breaks out in hives, the student's throat swells and tightens. Without immediate medical treatment the student could die within minutes.>> [Note to principal: This is a sample paragraph for a child with peanut or tree nut allergies. The same type of information will also be required if the student is allergic to fish such as tuna or some other product.]

All of our school staff, including lunch program staff, have bee made aware of this situation, and have been instructed in the correct procedures regarding anaphylactic shock by public health professionals. Prevention, of course, is the best approach for reducing the risk of a severe reaction, therefore we are requesting your cooperation in refraining from sending these food products, <<name of food products>>, to school with your child. [Note to principal: The following information may be included for student with peanut or nut allergies: "Even though the student is allergic to peanuts and peanut products, we have included tree nuts in our request because most nuts are processed with peanuts and are, therefore, contaminated with peanuts."]

We have told the other students of the problem, and have asked them not to share snacks, treats, or lunches with the severely allergic student. [Note to principal: Please see section 7 of Administrative Procedure 312 and ensure that parental consent has been obtained.] We have put procedures in place for snacks and the lunchroom, <<describe the procedures, separate table for student or separate table fro those with the peanut butter or tuna sandwiches as the case may be, or whatever approach you school has chosen to use>>. [Note to principal: It may be helpful to describe what the lunch program procedures may be in a general way.]

We realize that this request may post an inconvenience for you when preparing your student's snack and lunch, and we want to express our appreciation for your support and understanding about this severe allergy. In the near future, our school will be announcing a parent meeting for you to learn more about this situation. If you have any questions or concerns, please call me. Additional information is found in the Holy Spirit School Division Administrative Policy Manual which you may attain from the principal, or on our website at www.holyspirit.ab.ca.

On behalf of the student and his/her parents, I thank you for your co-operation.

Sincerely, The Principal

# **Anaphylaxis:**

Any delay in treatment could be fatal."

# Know what it is.

Anaphylaxis is a severe, life-threatening allergic reaction. It can be triggered by certain types of food (like peanuts and shellfish), insect stings, medicine, latex, exercise and unknown causes.

The following symptoms of anaphylaxis can occur within minutes or several hours after exposure to an allergy trigger:		
MOUTH: itching, swelling of the lips and/or tongue	THROAT*: itching, tightness, closure, hoarseness	
SKIN: itching, hives, redness, swelling	GUT: vomiting, diarrhea, cramps	
LUNG*: shortness of breath, cough, wheeze	HEART*: weak pulse, dizziness, fainting	

Only a few of these symptoms may be present.

\*Some symptoms can be life-threatening. ACT FAST!

# Know what to do.

Epinephrine (the active ingredient in the EpiPen Auto-Injector) is the medication recognized by healthcare professionals as the emergency treatment of choice for severe allergic reactions.

If any of the symptoms listed above are exhibited, administer the EpiPen\* Auto-Injector immediately.



- Hold firmly with orange tip pointing downward.
- Remove blue safety cap by pulling straight up. Do not bend or twist.



- Swing and push orange tip firmly into mid-outer thigh until you hear a 'click'.
- Hold on thigh for several seconds.



#### Built-in needle protection

 When the EpiPen- Auto-Injector is removed, the orange needle cover automatically extends to cover the injection needle, ensuring the needle is never exposed.



After administration, patients should seek medical attention immediately or go to the emergency room. For the next 48 hours, patients must stay within close proximity to a healthcare facility or where they can call 911.

For more information, or to order more posters, go to EpiPen.ca

EpiPen» and EpiPen» in Auto-Injectors are indicated for the emergency treatment of anaphylactic reactions in patients who are determined to be at increased risk for anaphylacts, including Individuals with a history of anaphylactic reactions

Tills product may not be right for you. Always much and below the product late

Beforences: 1. Liebertun P. Grop V. Opperheiter J. et al. The Garposts and management of auphylatios, an opticity location parameter. J Mercy Ots immunol 2005; 175:5483—52 2. Berns V. Locker VE. Senon IEE, on behalf of the World Lieve Organization author Connection on Connection in Association. More 2008-63: 1663—1693.







# **EpiPen**® Frequently asked questions

Retrieved from <a href="http://www.epipen.ca/en/about-epipen/faq">http://www.epipen.ca/en/about-epipen/faq</a> on May 1, 2015

# Q When should I inject EpiPen®?

Epinephrine should be administered as early as possible after the onset of symptoms of a severe allergic response. People requiring epinephrine will not always have predictable reactions. Adequate warning signs are not always present before serious reactions occur.

It is recommended that epinephrine be given at the start of any reaction associated with a known or suspected allergen contact. For people with a history of a severe cardiovascular collapse on exposure to an allergen, the physician may advise that epinephrine be administered immediately after exposure to that allergen, and before any reactionhas begun.

Epinephrine, when used as directed *immediately* following exposure to a known allergen, may prove life-saving.

### Q Why should I swing my arm to inject?

It is recommended that you swing your arm to inject EpiPen<sup>®</sup>. You do not need to apply excessive force to the swing, just enough to gain sufficient momentum to ensure the Auto-Injector deploys. Swinging your arm helps provide enough load on the tip of the EpiPen<sup>®</sup> to deliver an intramuscular (IM) injection. Administering the EpiPen<sup>®</sup> in this manner also helps facilitate the ideal angle for the IM injection into the anterolateral aspect of the thigh, which is perpendicular or at 90 degrees to the injection site.

# Q How many seconds should I hold EpiPen® in place when I inject?

The instructions on the EpiPen<sup>®</sup> cartridge state you should swing the arm and push the orange tip firmly into the mid-outer thigh so it "clicks" and hold it on the thigh for several seconds. By holding the EpiPen<sup>®</sup> in place at the injection site for several seconds, it helps assure that the dose of epinephrine has been delivered satisfactorily.

# Q How do I know if EpiPen® has been activated?

The extended orange needle cover and obscured window indicates that the EpiPen<sup>®</sup> has delivered the drug.

### Q How do I know if I have received my injection?

There is a viewing window on the barrel of the EpiPen<sup>®</sup> which prior to delivery is clear and after delivery is obscured or shaded. Also, once the dose is administered and the EpiPen<sup>®</sup> is released from the injection site, the orange needle cover extends and immediately locks in place encasing the needle to prevent accidental needle sticks. Both of these things serve as a visual indicator that the dose of epinephrine has been delivered.

Further, after the orange needle cover is extended the EpiPen<sup>®</sup> will not fit back into the carrier tube.

# Q Can I inject EpiPen® through my clothes?

Yes. If necessary, you can inject EpiPen® through clothing.

## Q What do I do in the event of an accidental injection into hands or feet, etc?

Accidental injection into the hand or feet may result in loss of blood flow to the affected areas and should be avoided. If there is an accidental injection into these areas, go immediately to the nearest emergency room for treatment.

# Q What do I do after I have used EpiPen®?

After using EpiPen<sup>®</sup>, seek medical attention immediately. Take the used EpiPen<sup>®</sup> with you to the nearest Emergency Department and give it to the Emergency Department staff for proper disposal. You can also bring any used EpiPen<sup>®</sup> Auto-Injectors to your pharmacy.

# Q How should I dispose of a used EpiPen®?

After using EpiPen<sup>®</sup> medical attention should be sought. The used EpiPen<sup>®</sup> should be taken to the nearest Emergency Room for proper disposal and this will also show the medical personnel what medication has already been administered.

For an expired EpiPen<sup>®</sup>, contact your local pharmacy, they should be able to assist in disposal of the product according to regulations.